

# An Overview of Low-Level Jets (LLJs) and Their Roles on Heavy Rainfall Over the Taiwan Area During the Early Summer Rainy Season

Yi-Leng Chen<sup>1</sup>, Chuan-Chi Tu<sup>2</sup>, Feng Hsiao<sup>1</sup>, Ching-Sen Chen<sup>2</sup>,  
Pay-Liam Lin<sup>2</sup>, and Po-Hsiung Lin<sup>3</sup>

<sup>1</sup>Department of Atmospheric Sciences, University of Hawai‘i at Mānoa, Honolulu, HI

<sup>2</sup>Department of Atmospheric Sciences, National Central University, Jhongli, Taiwan.

<sup>3</sup>Department of Atmospheric Sciences, National Taiwan University, Taipei, Taiwan

## ABSTRACT

During the early summer rainy season over Taiwan, three types of low-level jets are observed including a synoptic low-level jet (SLLJ), situated in the 850-hPa–700-hPa layer in the frontal zone, a marine boundary layer jet (MBLJ) embedded within the southwesterly monsoon flow over the northern South China Sea at approximately the 925-hPa level, and orographically induced jet at approximately the 1-km level off the northwestern Taiwan coast (e.g., barrier jet (BJ)). The characteristics and physical processes of the formation of these three types of low-level jets are reviewed, as well as and their roles responsible for the development of heavy rainfall are discussed.

\*Correspondence author address: Dr. Yi-Leng Chen, Department of Atmospheric Sciences, SOEST, University of Hawai‘i at Mānoa, 2525 Correa Road, Honolulu, HI 96822, USA  
Email: [yileng@hawaii.edu](mailto:yileng@hawaii.edu) Phone: (808) 956-2570 Fax: (808) 956-2877